



TITLE:

BROWN-PETERSON COHOMOLOGY OF $BP\mathbb{U}(p)$ (Cohomology Theory of Finite Groups and Related Topics)

AUTHOR(S):

KAMEKO, MASAKI; YAGITA, NOBUAKI

CITATION:

KAMEKO, MASAKI ...[et al]. BROWN-PETERSON COHOMOLOGY OF $BP\mathbb{U}(p)$ (Cohomology Theory of Finite Groups and Related Topics). 数理解析研究所講究録 2006, 1466: 103-106

ISSUE DATE:

2006-01

URL:

<http://hdl.handle.net/2433/48045>

RIGHT:

BROWN-PETERSON COHOMOLOGY OF $BPU(p)$

MASAKI KAMEKO (亀子正喜 富山国際大学地域学科)

NOBUAKI YAGITA (柳田伸顕 茨城大学教育学部)

Let p be a fixed odd prime and denote by $BP^*(X)$ (resp. $P(m)^*(X)$) the Brown-Peterson cohomology of a space X with the coefficient ring $BP^* = \mathbb{Z}_{(p)}[v_1, v_2, \dots]$ (resp. $P(m)^* = \mathbb{Z}/p[v_m, v_{m+1}, \dots]$) where $\deg v_k = -2p^k + 2$. We denote by $PU(n)$ the projective unitary group which is the quotient of the unitary group $U(n)$ by its center S^1 . Recall that the cohomologies of $PU(p)$ and exceptional Lie groups F_4, E_6, E_7, E_8 have odd torsion elements. In this paper, we compute the Brown-Peterson cohomologies of classifying spaces BG of these Lie groups G as BP^* -modules using the Adams spectral sequence. Let us write $H^*(X; \mathbb{Z}/p)$ by simply $H^*(X)$ and let \mathcal{A} be the mod p Steenrod algebra.

Our main result is as follows:

Theorem 0.1. *Let (G, p) be one of cases $(G = PU(p), p)$ for an arbitrary odd prime p and $G = F_4, E_7$ for $p = 3$, and $G = E_8$ for $p = 5$. Then the E_2 -terms of the Adams spectral sequences abutting to $BP^*(BG)$ and $P(m)^*(BG)$ for $m \geq 1$*

$$\text{Ext}_{\mathcal{A}}^{s,t}(H^*(BP), H^*(BG)), \quad \text{Ext}_{\mathcal{A}}^{s,t}(H^*(P(m)), H^*(BG))$$

have no odd degree elements.

As an immediate consequence is as follows:

Corollary 0.2. *For (G, p) in Theorem 1.1, the Adams spectral sequence abutting to $BP^*(BG)$ and $P(m)^*(BG)$ in the previous theorem collapse at the E_2 -level. In particular $BP^{\text{odd}}(BG) = P(m)^{\text{odd}}(BG) = 0$.*

Recall $K(m)^*(X) \cong K(m)^* \otimes_{P(m)^*} P(m)^*(X)$ is the Morava K -theory. From above theorem and corollary, we see $K(m)^{\text{odd}}(BPU(p)) = 0$. Then we have the following corollary ([Ko-Ya], [Ra-Wi-Ya])

Corollary 0.3. *For (G, p) in Theorem 1.1, the following holds:*

- (1) $BP^*(BG)$ is BP^* -flat for $BP^*(BP)$ -modules, i.e.,
 $BP^*(BG \times X) \cong BP^*(BG) \otimes_{BP^*} BP^*(X)$ for all finite complexes X
- (2) $K(n)^*(BG) \cong K(n)^* \otimes_{BP^*} BP^*(BG)$.
- (3) $P(n)^*(BG) \cong P(n)^* \otimes_{BP^*} BP^*(BG)$.

We give the BP^* -module structure of $BP^*(BPU(p))$ more explicitly, in this talk.

Theorem 0.4. *There is a BP^* -algebra isomorphism*

$$0 \rightarrow BP^* \hat{\otimes} M \rightarrow \text{gr} BP^*(BPU(p)) \rightarrow BP^* \hat{\otimes} IN / (f_0, f_1) \rightarrow 0$$

where

1. $M \cong \mathbb{Z}_{(p)}[x_4, x_6, \dots, x_{2p}]$ as $\mathbb{Z}_{(p)}$ -modules (but not $\mathbb{Z}_{(p)}$ -algebras).
2. $IN \cong \mathbb{Z}_{(p)}[x_{2p+2}, x_{2p(p-1)}] \{x_{2p+2}\}$; the principal ideal of $\mathbb{Z}[x_{2p+2}, x_{2p(p-1)}]$ generated by x_{2p+2} .

MASAKI KAMEKO (亀子正喜 富山国際大学地域学科) NOBUAKI YAGITA (柳田伸顕 茨城大学教育学部)

3. relations f_0, f_1 are given with modulo $(p, v_1, v_2, \dots)^2$

$$f_0 \equiv v_0 - v_2 x_{2p+2}^{p-1} + \dots, \quad f_1 \equiv v_1 - v_2 x_{2p(p-1)} + \dots$$

Remark 0.5. In the above theorem, suffix i of x_i means its degree. $BP^*(BPU(p))$ does not contain the subalgebra $BP^* \hat{\otimes} \mathbb{Z}_{(p)}[x_4, \dots, x_{2p}]$, but contains a subalgebra which is isomorphic as BP^* -modules to the above BP -subalgebra.

For an algebraic group G over \mathbb{C} , Totaro defines its Chow ring [To] and conjectures that $BP^*(BG) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong CH^*(BG)_{(p)}$. Recall that $PGL(p, \mathbb{C})$ is the algebraic group over \mathbb{C} corresponding the Lie group $PU(p)$.

Theorem 0.6. There is the isomorphism

$$BP^*(BPU(p)) \otimes_{BP^*} \mathbb{Z}_{(p)} \cong CH^*(BGL(p, \mathbb{C}))_{(p)}.$$

Hence there is the additive isomorphism

$$CH^*(BGL(p, \mathbb{C}))_{(p)} \cong \mathbb{Z}_{(p)}[x_4, x_6, \dots, x_{2p}] \oplus \mathbb{F}_p[x_{2p+2}, x_{2p(p-1)}]\{x_{2p+2}\}.$$

Remark. Recently Vistoli [Vi] also determined the additive structure of the Chow ring and integral cohomology of $BPGL(p, \mathbb{F}_p)$ by using stratified methods of Vessiozi. Moreover he shows that for $G = PGL(p, \mathbb{C})$

$$H^*(G; \mathbb{Z}) \rightarrow H^*(BT; \mathbb{Z})^{W_G(T)}$$

is epic.

Let $MGL^{*,*}(X)$ be the motivic cobordism ring defined by V.Voevodsky [Vo] and $MGL^{2*,*}(X) = \oplus_i MGL^{2i,i}(X)$.

Corollary 0.7. $MGL^{2*,*}(BPGL(p, \mathbb{C}))_{(p)} \cong MU^*(BPU(p))_{(p)}$.

We prove Theorem 1.1 using the Adams spectral sequence converging to the Brown-Peterson cohomology. The E_1 -term of the spectral sequence could be given by

$$\mathbb{F}_p[v_0, v_1, \dots] \hat{\otimes} H^*(X) \quad \text{with} \quad d_1 x = \sum_{k=0}^{\infty} v_k Q_k x$$

where Q_k 's are Milnor's operations. By the change-of-rings isomorphism, the E_2 -term is

$$\text{Ext}_{\mathcal{A}}(H^*(BP), H^*(X)) \cong \text{Ext}_{\mathcal{E}}(\mathbb{F}_p, H^*(X))$$

where $\mathcal{E} = \Lambda(Q_0, Q_1, \dots)$. The E_{∞} -term is given by $grBP^*(X)$.

To state the cohomology $H^*(BPU(p))$, we recall the Dickson algebra. Let A_n be an elementary abelian p -group of rank n , and

$$H^*(BA_n) \cong \mathbb{F}_p[t_1, \dots, t_n] \otimes \Lambda(dt_1, \dots, dt_n) \quad \text{with} \quad \beta(dt_i) = t_i.$$

The Dickson algebra is

$$D_n = \mathbb{F}_p[t_1, \dots, t_n]^{GL(n, \mathbb{F}_p)} \cong \mathbb{F}_p[c_{n,0}, \dots, c_{n,n-1}]$$

with $|c_{n,i}| = 2(p^n - p^i)$. The invariant ring under $SL(n, \mathbb{F}_p)$ is also given

$$SD_n = \mathbb{F}_p[t_1, \dots, t_n]^{SL(n, \mathbb{F}_p)} \cong D_n \{1, e_n, \dots, e_n^{p-2}\} \quad \text{with} \quad e_n^{p-1} = c_{n,0}.$$

We also recall the Mui's ([Mu]) result by using Q_i by [Ka-Mi]

$$grH^*(BA)^{SL(n, \mathbb{F}_p)} \cong SD_n/(e_n) \oplus SD_n \otimes \Lambda(Q_0, \dots, Q_{n-1})\{u_n\}$$

where $u_n = dt_1 \dots dt_n$ and $e_n = Q_0 \dots Q_{n-1} u_n$.

Theorem 0.8. *There is the short exact sequence*

$$0 \rightarrow M/p \rightarrow H^*(BP\mathbb{U}(p)) \rightarrow N \rightarrow 0$$

where M/p is the trivial \mathcal{E} -module given in Theorem 1.4 and

$$N = SD_2 \otimes \Lambda(Q_0, Q_1)\{u_2\} \cong \mathbb{F}_p[x_{2p+2}, x_{2(p^2-p)}] \otimes \Lambda(Q_0, Q_1)\{u_2\}$$

identifying $x_{2p+2} = e_2$ and $x_{2(p^2-p)} = c_{2,1}$.

This theorem is proved by using the following facts. The group $G = PU(p)$ has just two conjugacy classes of maximal elementary abelian p -subgroups, one of which is toral and the other is non-toral A of $\text{rank}_p = 2$. The cohomology $H^*(BG)$ is detected by this two subgroups. The restriction image to the non-toral subgroup is $i_A^*(H^*PU(p)) \cong H^*(BA)^{SL(2, \mathbb{F}_p)}$. Similar (but not same) facts also hold for the exceptional Lie groups given in Theorem 1.1.

Algebraic main result in this talk is as follows:

Theorem 0.9. *For $m \geq 0$, define f_0, \dots, f_{n-1} in $P(m)^* \hat{\otimes} SD_n$ by*

$$d_1 u_n = \sum_{k \geq m} v_k Q_k(u_n) = f_0 Q_0 u_n + \dots + f_{n-1} Q_{n-1} u_n.$$

Then the sequence f_0, \dots, f_{n-1} is a regular sequence in $P(m)^ \hat{\otimes} SD_n$.*

With the notation in this theorem, we prove that the complex

$$C = (P(m)^* \hat{\otimes} SD_n \otimes \Lambda(Q_0, Q_1, \dots, Q_{n-1})\{u_n\}, d_1)$$

with the differential $d_1 u_n = \sum_{i=0}^{n-1} f_i Q_i u_n$ is a Koszul complex. This means that

$$H_i(C, d_1) = \begin{cases} P(m)^* \hat{\otimes} SD_n \{e_n\} / (f_0, \dots, f_{n-1}) & \text{for } i = 0 \\ 0 & \text{for } i \geq 1. \end{cases}$$

Thus Theorem 1.1 follows from the above theorem.

Remark about the convergence of the Adams spectral sequence. By Theorem 15.6 in Boardman's paper [Bo2], since $H^*(BP)$ is of finite type, the above Adams spectral sequence is conditionally convergent. Moreover, since we prove the above Adams spectral sequence collapses at the E_2 -level, by the remark after Theorem 7.1 in [Bo1], the above Adams spectral sequence is strongly convergent, so that we know the Brown-Peterson cohomology up to group extension.

REFERENCES

- [A-G-M-V] K.Andersen, J.Grodal, J.Moller and A.Viruel, The classification of p -compact groups for p odd. *preprint arXiv:math. AT/0302346* v1 27 Feb. (2003).
- [Bo1] J.M.Boardman, Operations an the Adams spectral sequence for Brown-Peterson homology and cohomology. *preprint* (1988).
- [Bo2] J.M. Boardman, Conditionally Convergent Spectral Sequences. *Contemp. Math.*, **239** (1999) 49-84.
- [Ka-Mi] M.Kameko and M.Mimura, Mui invariant and Milnor operations *preprint*
- [Ka1] M.Kameko Poincare series of cotorsion products *Preprint*
- [Ka2] M.Kameko In preparation.
- [Ko-Mi] A.Kono and M.Minura, Cohomology of mod 3 of the classifying space of the Lie group E_6 . *Math. Scand.* **46** (1978), 223-235.
- [Ko-Ya] A. Kono and N. Yagita. Brown-Peterson and ordinary cohomology theories of classifying spaces for compact Lie groups. *Trans. of AMS.* **339** (1993), 781-798.
- [Mu] H.Mui. Modular invariant theory and the cohomology algebras of symmetric groups. *J.Fac.Sci.U. of Tokyo* **22** (1975), 319-369.

MASAKI KAMEKO (亀子正喜 富山国際大学地域学科) NOBUAKI YAGITA(柳田伸顕 茨城大学教育学部)

- [Toda] H.Toda Cohomology mod 3 of the classifying space BF_4 of the exceptional group F_4 . *J.Math.Kyoto Univ.* **13** (1973) 97-115.
- [To] B. Totaro. The Chow ring of classifying spaces. *Proc.of Symposia in Pure Math. "Algebraic K-theory" (1997:University of Washington,Seattle)* **67** (1999), 248-281.
- [Ve] G.Vezzosi. On the Chow ring of the classifying stack of $PGL_{3,C}$. *J.Reine Angew. Math.* **523** (2000), 1-54.
- [Vo] V. Voevodsky. The Milnor conjecture. *www.math.unic.edu/K-theory/0170* (1996)
- [Ra-Wi-Ya] D. C. Ravenel, W. S. Wilson and N. Yagita. Brown-Peterson cohomology from Morava K -theory. *K-theory* **15** (1998), 147-199.
- [Vi] A.Vistoli. On the cohomology and the chow ring of the classifying space of PGL_p . *http:hopf.math.purdue.edu* (2005), May.
- [Te-Ya] M. Tezuka and N.Yagita. The varieties of the mod p cohomology rings of extraspecial p -groups for an odd prime p . *Math. Proc. Cambridge Phil. Soc.* **94** (1983) 449-459.
- [Ya] N.Yagita. Applications of Atiyah-Hirzebruch spectral sequences for motivic cobordism. *Proc. London Math. Soc.* **90** (2005) 783-816.

Faculty of Regional science, Toyama University of Internal Studies, Toyama,
Japan

Faculty of Education, Ibaraki University, Mito, Ibaraki, Japan
kameko@tuins.ac.jp, yagita@mx.ibaraki.ac.jp